## Protect your community through regular **COVID-19 vaccinations**

A COVID-19 vaccination every 6-12 months is still the best way to protect yourself and others from severe illness, hospitalisation and death from COVID-19.

Unlike the flu and other respiratory viruses that peak in winter, COVID-19 has shown to spread year-round. New variants are continually emerging, even during warmer weather.

Frequency of COVID-19 vaccination	Recommended	Eligible
People aged 75 years and older	Every 6 months	Every 6 months
People aged 65 to 74 years People aged 18 -74 with severe immunocompromise	Every 12 months	Every 6 months
People aged 18 – 64 without severe immunocompromise	Every 12 months	Every 12 months

\*If it is not known when the last COVID-19 vaccine or COVID-19 infection occurred, it is still safe to get another COVID-19 vaccine.

Schedule **COVID-19** vaccination reminders, obtain patient Maintain summer consent, and plan for co-administration with flu vaccine during yearly health checks.

Schedule COVID-19 vaccination reminders for eligible patients.

\*Puno! (Jun-Jul)

Stay alert to upper respiratory infections like COVID-19 to reduce spread in spring.

vaccinations.

Kambarang

Plan COVID-19 and flu vaccine co-administration.

Protect the community with **COVID-19** vaccination and flu co-administration.



